

## Congenital Toxoplasmosis in a Indo-Pacific Bottlenose Dolphin (*Tursiops aduncus*)

**J. E. Jardine and J. P. Dubey**\*†, VETPATH Laboratory Services, P.O. Box 18, Belmont 6984, Western Australia; and \*Parasite Biology, Epidemiology and Systematics Laboratory, Building 1001, Animal Natural Resources Institute, Agricultural Research Service, U.S. Department of Agriculture, Beltsville Agricultural Research Center-East, Beltsville, Maryland 20705-2350. †To whom correspondence should be addressed *e-mail*: [jdubey@anri.barc.usda.gov](mailto:jdubey@anri.barc.usda.gov)

**ABSTRACT:** *Toxoplasma gondii* was identified in tissues of a stillborn late-term fetus from an Indo-Pacific bottlenose dolphin (*Tursiops aduncus*). Myocardial necrosis and nonsuppurative inflammation in the heart and nonsuppurative necrotizing encephalitis were associated with tachyzoites and tissue cysts. The diagnosis was confirmed by immunohistochemical staining with anti-*T. gondii*-specific polyclonal rabbit serum.

There are several reports of *Toxoplasma gondii*-associated disease in sea mammals (Van Pelt and Dieterich, 1973; Migaki et al., 1977, 1990; Holshuh et al., 1985; Mikaelian et al., 2000). The first report of toxoplasmosis in a dolphin was from Brazil (Bandoli and Oliveira, 1977). Cruickshank et al. (1990) and Inskeep et al. (1990) reported toxoplasmosis in Atlantic bottlenose dolphins (*Tursiops truncatus*) from North America, and Domingo et al. (1992) reported *T. gondii* infection from striped dolphins (*Stenella coeruleoalba*) from Spain. These reports were based on histologic evaluation of dead animals. Cole et al. (2000) isolated several strains of *T. gondii* from the brains of California sea lions (*Zalophus californianus*), and Miller, Crosbie et al. (2001) and Miller, Sverlow et al. (2001) isolated *T. gondii* from a Pacific harbor seal (*Phoca vitulina richardsi*), also from California. In addition, encephalitis associated with the related protozoon *Sarcocystis neurona* has been diagnosed in 3 sea otters (*Enhydra lutris*) (Rosonke et al., 1999; Lindsay et al., 2000; Miller, Crosbie et al., 2001) and 7 Pacific harbor seals (Lapointe et al., 1998). Dual *S. neurona* and *T. gondii* infections have been confirmed in a sea otter (Lindsay et al., 2001) and a Pacific harbor seal (Miller, Sverlow et al., 2001). Here, we report a transplacentally

transmitted *T. gondii* infection in a stillborn Indo-Pacific bottlenose dolphin (*Tursiops aduncus*) fetus from Australia.

A late-term fetus was born dead to a mature captive-born Indo-Pacific bottlenose dolphin in a large open-water enclosure and was retrieved after 3 days. The fetus weighed 10 kg (normal size is approximately 30 kg), and samples of brain, heart, lung, liver, kidney, lymph node, and gastrointestinal tract were collected during necropsy and preserved in buffered neutral 10% formalin for histological assessment. Paraffin-embedded sections were cut at 5 µm thickness, stained with hematoxylin and eosin (H&E), and examined microscopically. Selected sections were immunohistochemically stained with antibodies to *T. gondii* (Lindsay and Dubey, 1989) and *S. neurona* (Dubey and Hamir, 2000) as previously described.

Multiple tissues of the dolphin showed marked postmortem changes, including prominent invasion of the tissues with putrefactive bacteria. Important histological findings were recognized within the heart and the brain. The myocardium had multiple small areas of necrosis characterized by myofiber fragmentation and lysis with some mild nonsuppurative inflammation (Fig. 1). Protozoa were seen within the cells of the surrounding tissues. Some mild interstitial inflammation was evident in the epicardium. Similar protozoa were randomly distributed within the brain and were accompanied by a multifocal, nonsuppurative necrotizing encephalitis. No other important changes could be identified in the other tissues because of severe autolysis.

Tissue cysts and tachyzoites were both identified in close association with lesions (Fig. 1). Tissue cysts were often at the periphery of the lesion. The organisms stained strongly positive with anti-*T. gondii* antibodies and more organisms were seen in these sections than in those stained with H&E (compare Fig. 1A, B).




FIGURE 1. Lesions and *Toxoplasma gondii* in the brain (A–C) and the heart (D) of the dolphin fetus. A, B. Focal lesion of necrosis (n), inflammation, and a group of protozoa (arrows). C. Higher magnification of the *T. gondii* (arrow) in Figure 1A. Note partly degenerated tachyzoites (arrows) and a small tissue cyst (arrowhead). Bacillary rods are also present. D. Focal necrosis (n) and groups of *T. gondii* (arrows). H&E (A, C) and immunohistochemical stain with anti-*T. gondii* antibodies (B, D).

Although toxoplasmosis has been described in several species of marine animals, this is the first report of congenital toxoplasmosis in a marine animal. The dam of the fetus lived in a netted enclosure within a public marina and probably acquired the *T. gondii* infection by ingestion of food or water contaminated with oocysts. Feral cats frequent the quayside area.

#### LITERATURE CITED

- BANDOLI, J. G., AND C. A. B. DE OLIVEIRA. 1977. Toxoplasmose em *Sotalia guianensis* (van Beneden, 1863), Cetacea—Delphinidae. *Folha Médica* 75: 459–468.
- COLE, R. A., D. S. LINDSAY, D. K. HOWE, C. L. RODERICK, J. P. DUBEY, N. J. THOMAS, AND L. A. BAETEN. 2000. Biological and molecular

- characterizations of *Toxoplasma gondii* strains obtained from southern sea otters (*Enhydra lutris nereis*). *Journal of Parasitology* **86**: 526–530.
- CRUICKSHANK, J. J., D. M. HAINES, N. C. PALMER, AND D. J. STAUBIN. 1990. Cysts of a *Toxoplasma*-like organisms in an Atlantic bottlenose dolphin. *Canadian Veterinary Journal* **31**: 213–215.
- DOMINGO, M., J. VISA, M. PUMAROLA, A. J. MARCO, L. FERRER, R. RABANAL, AND S. KENNEDY. 1992. Pathologic and immunocytochemical studies of morbillivirus infection in striped dolphins (*Stenella coeruleoalba*). *Veterinary Pathology* **29**: 1–10.
- DUBEY, J. P., AND A. N. HAMIR. 2000. Immunohistochemical confirmation of *Sarcocystis neurona* infections in raccoons, mink, cat, skunk, and pony. *Journal of Parasitology* **86**: 1150–1152.
- HOLSHUH, H. J., A. E. SHERROD, C. R. TAYLOR, B. F. ANDREWS, AND E. B. HOWARD. 1985. Toxoplasmosis in a feral northern fur seal. *Journal of the American Veterinary Medical Association* **187**: 1229–1230.
- INSKEEP, I. W., C. H. GARDINER, R. K. HARRIS, J. P. DUBEY, AND R. T. GOLDSTON. 1990. Toxoplasmosis in Atlantic bottle-nosed dolphins (*Tursiops truncatus*). *Journal of Wildlife Diseases* **26**: 377–382.
- LAPOINTE, J. M., P. J. DUIGNAN, A. E. MARSH, F. M. GULLAND, B. C. BARR, D. K. NAYDAN, D. P. KING, C. A. FARMAN, K. A. B. HUNTINGDON, AND L. J. LOWENSTINE. 1998. Meningoencephalitis due to a *Sarcocystis neurona*-like protozoan in Pacific harbor seals (*Phoca vitulina richardsi*). *Journal of Parasitology* **84**: 1184–1189.
- LINDSAY, D. S., AND J. P. DUBEY. 1989. Immunohistochemical diagnosis of *Neospora caninum* in tissue sections. *American Journal of Veterinary Research* **50**: 1981–1983.
- , N. J. THOMAS, AND J. P. DUBEY. 2000. Biological characterization of *Sarcocystis neurona* from a southern sea otter (*Enhydra lutris nereis*). *International Journal for Parasitology* **30**: 617–624.
- , ———, A. C. ROSYPAL, AND J. P. DUBEY. 2001. Dual *Sarcocystis neurona* and *Toxoplasma gondii* infection in a northern sea otter from Washington state, USA. *Veterinary Parasitology* **97**: 319–327.
- MIGAKI, G., J. F. ALLEN, AND H. W. CASEY. 1977. Toxoplasmosis in a California sea lion (*Zalophus californianus*). *American Journal of Veterinary Research* **38**: 135–136.
- , T. R. SAWA, AND J. P. DUBEY. 1990. Fatal disseminated toxoplasmosis in a spinner dolphin (*Stenella longirostris*). *Veterinary Pathology* **27**: 463–464.
- MIKHAELIAN, I., J. BOISCLAIR, J. P. DUBEY, S. KENNEDY, AND D. MARTINEAU. 2000. Toxoplasmosis in beluga whales (*Delphinapterus leucas*) from the St. Lawrence estuary: Two case reports and a serologic survey. *Journal of Comparative Pathology* **122**: 73–76.
- MILLER, M. A., P. R. CROSBIE, K. SVERLOW, K. HANNI, B. C. BARR, N. KOCK, M. J. MURRAY, L. J. LOWENSTINE, AND P. A. CONRAD. 2001. Isolation and characterization of *Sarcocystis* from brain tissue of a free-living southern sea otter (*Enhydra lutris nereis*) with fatal meningoencephalitis. *Parasitology Research* **87**: 252–257.
- , K. SVERLOW, P. R. CROSBIE, B. C. BARR, L. J. LOWENSTINE, F. M. GULLAND, A. PACKHAM, AND P. A. CONRAD. 2001. Isolation and characterization of two parasitic protozoa from a Pacific harbor seal (*Phoca vitulina richardsi*) with meningoencephalitis. *Journal of Parasitology* **87**: 816–822.
- ROSONKE, B. J., S. R. BROWN, S. J. TORNQUIST, S. P. SNYDER, M. M. GARNER, AND L. L. BLYTHE. 1999. Encephalomyelitis associated with a *Sarcocystis neurona*-like organism in a sea otter. *Journal of the American Veterinary Medical Association* **215**: 1839–1842.
- VAN PELT, R. W., AND R. A. DIETERICH. 1973. Staphylococcal infection and toxoplasmosis in a young harbor seal. *Journal of Wildlife Diseases* **9**: 258–261.